

The Three-step Synthesis of 2,6-Dinitro-4-methylaniline

Padao Pha
Saint Catherine University

Stefani Ly
Saint Catherine University

Follow this and additional works at: http://sophia.stkate.edu/undergraduate_research_symposium

Pha, Padao and Ly, Stefani, "The Three-step Synthesis of 2,6-Dinitro-4-methylaniline" (2013). *Sr. Seraphim Gibbons Undergraduate Symposium*. 7.

http://sophia.stkate.edu/undergraduate_research_symposium/2013/Sciences/7

This Event is brought to you for free and open access by the Conferences and Events at SOPHIA. It has been accepted for inclusion in Sr. Seraphim Gibbons Undergraduate Symposium by an authorized administrator of SOPHIA. For more information, please contact hltompkins@stkate.edu.

The Three-step Synthesis of 2,6-Dinitro-4-methylaniline

Padao Pha,
Stephani Ly

This experiment entails the three-step synthesis of 2,6-Dinitro-4-methylaniline from *p*-toluidine. *P*-toluidine was tosylated using tosyl chloride. The subsequent product was then dinitrated with nitric acid and then sulfonated using sulfuric acid to yield 2,6-dinitro-4-methylaniline. While *p*-toluidine is known as a suspected cancer agent, 2,6-Dinitro-4-methylaniline is believed to have possible medicinal purposes making it important for future health applications. This synthesis was made more green by replacing sodium hydroxide with ethanol in step one. This is “greener” because ethanol is less corrosive and less health hazardous.