

# Synthesis of a Derivative for Metal Salen

Hlee Vang  
*St. Catherine University*

Tippaket Thammavong  
*St. Catherine University*

Follow this and additional works at: [https://sophia.stkate.edu/undergraduate\\_research\\_symposium](https://sophia.stkate.edu/undergraduate_research_symposium)

---

Vang, Hlee and Thammavong, Tippaket, "Synthesis of a Derivative for Metal Salen" (2013). *Sr. Seraphim Gibbons Undergraduate Symposium*. 33.

[https://sophia.stkate.edu/undergraduate\\_research\\_symposium/2013/Sciences/33](https://sophia.stkate.edu/undergraduate_research_symposium/2013/Sciences/33)

This Event is brought to you for free and open access by the Conferences and Events at SOPHIA. It has been accepted for inclusion in Sr. Seraphim Gibbons Undergraduate Symposium by an authorized administrator of SOPHIA. For more information, please contact [amshaw@stkate.edu](mailto:amshaw@stkate.edu).

## Synthesis of a Derivative for Metal Salen

Hlee Vang

Tippaket Thammavong

### Abstract

This three step procedure outlines the synthesis of 5-(2-sulfhydrylethyl)salicylaldehyde.

First the intermediate, 4-(2-iodoethyl)phenol ~~is was~~ made by combining hydriodic acid and 4-methoxyphenethyl. ~~Then we treat the~~The iodinated product was converted into a intermediate~~with a~~ Grignard reagent and treated with; paraformaldehyde; and triethylamine to make 5-(2-iodoethyl)salicylaldehyde. Lastly, 5-(2-sulfhydrylethyl)salicylaldehyde was obtained by reacting 5-(2-iodoethyl)salicylaldehyde with thiourea. Benzene is a known carcinogenic compound. Here, it is replaced by toluene because it has similar physical properties but is less carcinogenic. The product of this synthesis can be used to form derivatives of metal salens which could play a valuable role in medical technology and engineering.